

An Extension of Phenomenological Renormalization Method

M. Itakura

Department of Pure and Applied Sciences,

University of Tokyo,

Meguro-ku, Komaba 3-8-1, Tokyo 153, Japan

Abstract

We present an extension of the so-called cumulant crossing method which is used for determination of critical point in Monte Carlo simulations. The new method uses linear combination of several different order-parameter moments and almost eliminates the systematic deviation of crossing points from the true critical point. The performance of the method is tested by applying it to the 2D Ising model.

PACS numbers: 68.35.Rh, 52.65.Pp

The phenomenological renormalization method [1] has been widely used in Monte-Carlo simulations of critical phenomena. It make use of the property of reduced order-parameter moments such as

$$U_L^{(n)} \equiv \langle M^n \rangle_L / \langle M^2 \rangle_L^{n/2} \quad (1)$$

where M is the order-parameter and $\langle \dots \rangle_L$ denotes the thermal average in finite system with linear size L . In the vicinity of the critical point, they obey following scaling form [1]:

$$U_L^{(n)} \sim f_{n,0}(tL^{1/\nu}) + L^{-\omega_1} f_{n,1}(tL^{1/\nu}) + L^{-\omega_2} f_{n,2}(tL^{1/\nu}) + \dots \quad (2)$$

where t is the temperature deviation from the critical point and ν is the critical exponent related to the correlation length. The correction-to-scaling and correction-to-correction-to-scaling exponent etc. [2] are denoted by ω_i ($0 < \omega_1 < \omega_2 < \dots$) and $f_{n,i}$ denotes associated scaling functions. Two quantities, $U_L^{(n)}$ and $U_{bL}^{(n)}$, intersect at the following temperature:

$$t \sim C_1 L^{-1/\nu-\omega_1} + C_2 L^{-1/\nu-\omega_2} + \dots \quad (3)$$

$$C_i \sim \frac{(1 - b^{-\omega_i}) f_{n,i}(0)}{(b^{1/\nu} - 1) f'_{n,0}(0)} \quad (4)$$

One can extrapolate the critical point by plotting the crossing temperature versus $L^{-1/\nu-\omega_1}$. However, this method requires a knowledge about the exponents ν and ω_1 before the estimation of the critical point.

We present an extended method which reduces the systematic deviation of the crossing points from the true critical point and does not require any finite-size scaling. The key idea is that one can eliminate the correction-to-scaling terms by using linear combination of several different moments.

$$V_L \equiv a_4 U_L^{(4)} + a_6 U_L^{(6)} + \dots \quad (5)$$

It obeys the following scaling form:

$$V_L \sim F_0(tL^{1/\nu}) + L^{-\omega_1} F_1(tL^{1/\nu}) + L^{-\omega_2} F_2(tL^{1/\nu}) + \dots \quad (6)$$

where $F_i(x) \equiv a_4 f_{4,i}(x) + a_6 f_{6,i}(x) + \dots$ is the scaling function of V_L . One can adjust the coefficients a_n so that equations $F_1(0) = 0, F_2(0) = 0, \dots$ holds, which leads to $C_1 = 0, C_2 = 0, \dots$ in the equation(3). For example, consider the following simple case:

$$V_L = a_4 U_L^{(4)} + a_6 U_L^{(6)} \quad (7)$$

The systematic deviation can be reduced by adjusting the coefficients a_n ; the deviation is approximately proportional to:

$$\frac{a_4 f_{4,1}(0) + a_6 f_{6,1}(0)}{a_4 f'_{4,0}(0) + a_6 f'_{6,0}(0)} L^{-1/\nu-\omega_1} \quad (8)$$

Moreover, one can control the direction of the deviation by adjusting the coefficients a_n and determine the upper and lower bound of the critical point.

However, the estimation of appropriate coefficients a_n is a non-trivial part of this method.

We show an example of the way of estimation in the following.

If we plot $U_L^{(4)}$ versus $U_L^{(6)}$ using the values at the critical point, all points are approximately on a straight line.

$$(U_L^{(4)}(0), U_L^{(6)}(0)) \sim (f_{4,0}(0), f_{6,0}(0)) + L^{-\omega_1} (f_{4,1}(0), f_{6,1}(0)) \quad (9)$$

The slope of the line gives the ratio of two constants $f_{4,1}(0)$ and $f_{6,1}(0)$. The slope at the critical point can be approximated by the value at a temperature where the following condition is satisfied:

$$\frac{U_{bL}^{(4)} - U_L^{(4)}}{U_{bL}^{(6)} - U_L^{(6)}} = \frac{U_{cL}^{(4)} - U_L^{(4)}}{U_{cL}^{(6)} - U_L^{(6)}} \quad (10)$$

where $b \neq c \neq 1$ is some constant.

We have applied this method to Monte Carlo data of the 2D Ising model. Simulations were done at the critical coupling ($K_c = \log(1 + \sqrt{2})$) using Swendsen-Wang algorithm [3] and 10^5 observations were made after thermalization. Positions of crossing points were calculated with reweighting techniques [4]. Statistical errors were calculated with jackknife procedure.

Order parameter moments were calculated using cluster size distribution via the following equations:

$$\langle M^6 \rangle_L = 16 \langle S_6 \rangle_L - 30 \langle S_4 \rangle_L \langle S_2 \rangle_L + 15 \langle S_2 \rangle_L^3 \quad (11)$$

$$\langle M^4 \rangle_L = -2 \langle S_4 \rangle_L + 3 \langle S_2 \rangle_L^2 \quad (12)$$

$$\langle M^2 \rangle_L = \langle S_2 \rangle_L \quad (13)$$

$$S_n \equiv L^{-2n} \sum_j m_j^n \quad (14)$$

where m_j is the size of j 'th cluster in the cluster generation step of Swendsen-Wang algorithm.

Fig.1 shows the size-dependence of the deviations of crossing points in the two cases of using $U_L^{(4)}$ and $V_L = 5U_L^{(4)} + U_L^{(6)}$ with $b = 1/2$. One can see a significant reduction of deviation by the use of V_L . The residual deviations of the case of V_L may be the effect of higher order corrections. Further extension for eliminating higher order corrections is straightforward, but it needs observation of higher order moments of the order parameter, which is more subjected to statistical errors.

This work was supported by a Grant-in-Aid for Scientific Research by the Ministry of Education, Science and Culture.

REFERENCES

- [1] K. Binder, Z. Phys. **B 43**, 119 (1981).
- [2] F. J. Wegner, Phys. Rev. **B 5**, 4529 (1972).
- [3] R. H. Swendsen and J. S. Wang, Phys. Rev. Lett. **58**, 86 (1987).
- [4] A. M. Ferrenberg and R. H. Swendsen, Phys. Rev. Lett. **61**, 2635 (1988).

Figure Caption

Fig.1. Plot of crossing points in the case of using conventional ($U_L^{(4)}$) and new (V_L) parameter.

